

United States Department of Agriculture
Center for Veterinary Biologics
Testing Protocol

SAM 306

Supplemental Assay Method for the Titration of Feline
Calicivirus in Cell Culture

Date: April 4, 2005
Number: VIRSAM0306.03
Supersedes: MVSAM0306.02, Dated March 6, 2001
Standard Requirement: 9 CFR, Part 113.314
Contact: Victor M. Becerra, (515) 663-7468
Marsha J. Hegland, (515) 663-7659
Approvals:

/s/Donna M. Gatewood Date: 22APR05
Donna M. Gatewood, Section Leader
Virology Section

/s/Richard E. Hill, Jr. Date: 26APR05
Richard E. Hill, Jr., Director
Policy, Evaluation, and Licensing
Center for Veterinary Biologics

/s/Rebecca L.W. Hyde Date: 26APR05
Rebecca L.W. Hyde, Section Leader
Quality Assurance
Center for Veterinary Biologics

United States Department of Agriculture
Animal and Plant Health Inspection Service
P. O. Box 844
Ames, IA 50010

Mention of trademark or proprietary product does not constitute a guarantee or warranty of the product by USDA and does not imply its approval to the exclusion of other products that may be suitable.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

Table of Contents

1. Introduction
2. Materials
 - 2.1 Equipment/instrumentation
 - 2.2 Reagents/supplies
3. Preparation for the test
 - 3.1 Personnel qualifications/training
 - 3.2 Preparation of equipment/instrumentation
 - 3.3 Preparation of reagents/control procedures
 - 3.4 Preparation of the Test Vaccine
4. Performance of the test
5. Interpretation of the test results
6. Report of test results
7. References
8. Summary of revisions

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

1. Introduction

This Supplemental Assay Method (SAM) describes an *in vitro* titration method for assaying modified-live feline calicivirus (FCV) vaccines for viral content. The method uses plaque forming units (PFU) in a cell culture system for titration of FCV.

2. Materials

2.1 Equipment/instrumentation

Equivalent equipment or instrumentation may be substituted for any brand name listed below.

- 2.1.1 Syringe, self-refilling, repetitive, 2-ml
- 2.1.2 Micropipettor, 200- μ l, and tips (Rainin Pipetman®)
- 2.1.3 Blender
- 2.1.4 Media bottle, borosilicate glass with screw-top lid, 1000-ml
- 2.1.5 Incubator, $36^{\circ} \pm 2^{\circ}\text{C}$, high humidity, $5\% \pm 1\%$ CO_2 (Model 3158, Forma Scientific Inc.)
- 2.1.6 Water bath
- 2.1.7 Inverted light microscope (Model CK, Olympus America Inc.)
- 2.1.8 Vortex mixer (Vortex-2 Genie, Model G-560, Scientific Industries Inc.)
- 2.1.9 Pipette-aid

2.2 Reagents/supplies

Equivalent reagents or supplies may be substituted for any brand name listed below. All reagents and supplies must be sterile.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

2.2.1 FCV Reference, C-14 strain [available from the Center for Veterinary Biologics (CVB)]

2.2.2 Crandall feline kidney (CRFK) cell culture, free of extraneous agents as tested by the Code of Federal Regulations, Title 9 (9 CFR).

2.2.3 Minimum Essential Medium (MEM)

2.2.3.1 9.61 g MEM with Earles salts without bicarbonate

2.2.3.2 1.1 sodium bicarbonate (NaHCO_3)

2.2.3.3 Dissolve with 900 ml deionized water (DW).

2.2.3.4 Add 5.0 g lactalbumin hydrolysate or edamine to 10 ml DW. Heat to $60^\circ \pm 2^\circ\text{C}$ until dissolved. Add to **Section 2.2.3.3** with constant mixing.

2.2.3.5 Q.S. to 1000 ml with DW; adjust pH to 6.8-6.9 with 2N hydrochloric acid (HCl).

2.2.3.6 Sterilize through a 0.22- μm filter.

2.2.3.7 Aseptically add 50 $\mu\text{g/ml}$ gentamicin sulfate

2.2.3.8 Store at $2^\circ - 7^\circ\text{C}$.

2.2.4 Growth Medium

2.2.4.1 900 ml of MEM

2.2.4.2 Aseptically add:

1. 100 ml gamma-irradiated fetal bovine serum (FBS)

2. 10 ml L-glutamine (200 mM)

2.2.4.3 Store at $2^\circ - 7^\circ\text{C}$.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

2.2.5 2X Medium

2.2.5.1 19.22 g MEM with Earles salts without bicarbonate

2.2.5.2 2.2 g NaHCO_3

2.2.5.3 Dissolve with 900 ml DW.

2.2.5.4 Add 5.0 g lactalbumin hydrolysate or edamine to 10 ml DW. Heat to $60^\circ \pm 2^\circ\text{C}$ until dissolved. Add to **Section 2.2.5.3** with constant mixing.

2.2.5.5 Q.S. to 1000 ml with DW and adjust pH to 6.8-6.9 with 2N HCl.

2.2.5.6 Sterilize through a 0.22- μm filter.

2.2.5.7 Store at 2° - 7°C .

2.2.5.8 On the day of test initiation and prior to preparing the Overlay Medium, aseptically add 100 $\mu\text{g/ml}$ gentamicin sulfate.

2.2.6 2% Tragacanth Gum (Trag)

2.2.6.1 20 g Trag

2.2.6.2 1000 ml DW

2.2.6.3 Mix small amounts of Trag at a time vigorously, with a blender set on high.

2.2.6.4 Pour 500 ml each into 1000-ml media bottles.

2.2.6.5 Sterilize by autoclaving at 15 psi, $121^\circ \pm 2^\circ\text{C}$ for 35 ± 5 minutes.

2.2.6.6 Store at 2° - 7°C .

2.2.7 7.5% Sodium Bicarbonate

2.2.7.1 7.5 g NaHCO_3

2.2.7.2 Q.S. to 100 ml with DW.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

2.2.7.3 Sterilize by autoclaving at 15 psi, 121°± 2°C for 30 ± 10 minutes.

2.2.7.4 Store at 2°- 7°C.

2.2.8 70% Ethyl Alcohol

2.2.8.1 73 ml ethyl alcohol

2.2.8.2 27 ml DW

2.2.8.3 Store at room temperature.

2.2.9 Crystal Violet Stain

2.2.9.1 7.5 g crystal violet

2.2.9.2 50 ml 70% Ethyl Alcohol

2.2.9.3 Dissolve crystal violet in alcohol; add remaining ingredients.

2.2.9.4 250 ml formaldehyde

2.2.9.5 Q.S. to 1000 ml with DW.

2.2.9.6 Filter through Whatman® #1 filter paper.

2.2.9.7 Store at room temperature.

2.2.10 Tissue culture plates, 4-well

2.2.11 Polystyrene tubes, 12 x 75-mm

2.2.12 Pipettes, 25-ml

2.2.13 Needles, 18-gauge x 1 1/2-inch

2.2.14 Syringes (tuberculin), 1-ml

3. Preparation for the test

3.1 Personnel qualifications/training

Personnel must have experience in the preparation and maintenance of cell culture as well as in the propagation and maintenance of animal viruses and the quantitation of virus infectivity by plaque formation.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

3.2 Preparation of equipment/instrumentation

On the day of test initiation, set a water bath at $36^{\circ}\pm 2^{\circ}\text{C}$.

3.3 Preparation of reagents/control procedures

3.3.1 Preparation of CRFK tissue culture plates (Test Plates)

Cells are prepared from healthy, confluent CRFK cell cultures that are maintained by passing every 3 to 4 days. Two days prior to test initiation, add 8.0 ml/well of approximately $10^{5.0}$ to $10^{5.3}$ cells/ml suspended in Growth Medium into all wells of a 4-well tissue culture plate. Prepare 1 Test Plate for the FCV Reference Control and 1 for each Test Vaccine. Incubate at $36^{\circ}\pm 2^{\circ}\text{C}$ in a CO_2 incubator.

3.3.2 Preparation of FCV Reference Control

3.3.2.1 On the day of test initiation, rapidly thaw a vial of FCV Reference in the water bath.

3.3.2.2 Dispense 1.8 ml of MEM into sufficient 12 x 75-mm polystyrene tubes to bracket the expected endpoint according to the Center for Veterinary Biologics (CVB) Reagent Data Sheet; appropriately label (e.g., 9 tubes, labeled 10^{-1} through 10^{-9} , respectively).

3.3.2.3 Transfer 200 μl of the FCV Reference to the tube labeled 10^{-1} ; mix by vortexing.

3.3.2.4 Using a new pipette tip, transfer 200 μl from the 10^{-1} -labeled tube to the 10^{-2} tube; mix by vortexing.

3.3.2.5 Repeat **Section 3.3.2.4** for each of the subsequent dilutions, transferring 200 μl of the previous dilution to the next dilution tube until the tenfold dilution series is completed.

3.3.3 Overlay Medium Preparation

On the day of test initiation, prepare approximately 35 ml per plate of Overlay Medium.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

3.3.3.1 Aseptically add to 1 volume of 2X Medium:

1. 10% gamma-irradiated FBS
2. 2% 7.5% Sodium Bicarbonate
3. 1 volume of 2% Trag

3.3.3.2 Mix and warm the Overlay Medium in the water bath for 60 ± 10 minutes prior to performing the procedure in **Section 4.5**.

3.4 Preparation of the Test Vaccine

3.4.1 The initial test will be with a single vial (a single sample from 1 vial). On the day of test initiation, using a 1.0-ml syringe and an 18-gauge x 1 1/2-inch needle, rehydrate a vial of the Test Vaccine with the provided diluent by transferring 1.0 ml for a 1-ml-dose vaccine, 0.5 ml for a 1/2-ml-dose vaccine, etc., into the vial containing the lyophilized Test Vaccine; mix by vortexing. Incubate for 15 ± 5 minutes at room temperature.

3.4.2 Dispense 1.8 ml MEM into each of 6, 12 x 75-mm polystyrene tubes labeled 10^{-1} through 10^{-6} .

3.4.3 Transfer 200 μ l of the Test Vaccine to the tube labeled 10^{-1} ; mix by vortexing.

3.4.4 Using a new pipette tip, transfer 200 μ l from the tube labeled 10^{-1} to the 10^{-2} tube; mix by vortexing.

3.4.5 Repeat **Section 3.4.4** for each of the subsequent dilutions, transferring 200 μ l of the previous dilution to the next dilution tube until the tenfold dilution series is completed.

4. Performance of the test

4.1 On the day of test initiation, label the Test Plates and remove the Growth Medium.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

4.2 Inoculate 1 well/dilution with 200 μ l from dilutions 10^{-6} through 10^{-3} of the Test Vaccine, starting with the highest dilution (most dilute). Inoculate 1 well/dilution from dilutions 10^{-9} through 10^{-7} of the FCV Reference Control. Change tips between each unique sample (i.e., each Test Vaccine), but tip changes are not necessary between each dilution in a series when pipetting from the most dilute to the most concentrated (e.g., 10^{-9} through 10^{-7}). Gently rotate the plates to evenly disperse the inoculum.

4.3 One uninoculated well on a Test Plate serves as a negative cell control.

4.4 Incubate the inoculated Test Plates at $36^{\circ} \pm 2^{\circ}\text{C}$ for 75 ± 15 minutes in a CO_2 incubator. To evenly disperse the inoculum, gently rotate the plates at intervals of 25 ± 5 minutes.

4.5 Add 8 ml/well of the Overlay Medium to the plates with a 25-ml pipette. Discard any unused, warmed Overlay Medium.

4.6 Incubate the Test Plates undisturbed at $36^{\circ} \pm 2^{\circ}\text{C}$ in a CO_2 incubator for 4 days ± 1 day.

4.7 After incubation, without removing the Overlay Medium, pipette 5 ml of the Crystal Violet Stain (**Section 2.2.9**) into each well of the Test Plates with a 25-ml pipette; mix by gentle rotation.

4.8 Allow Test Plates to incubate at room temperature 25 ± 5 minutes.

4.9 Wash the Overlay Medium and the Crystal Violet Stain from the cell monolayers by dipping each plate several times in a container overflowing with running water from the cold water tap until the water is clear. Allow to air dry.

4.10 Count the number of FCV plaques for each well. If FRV and FCV plaques are counted together in a combination vaccine, the FCV plaques will contrast markedly from FRV. The FRV plaques are visible as small, clear plaques approximately 1 mm in diameter with distinct edges. The FCV plaques are large, clear circular areas (averaging 3 to 4 mm in diameter) with fuzzy edges. If the plaques are not distinguishable by size, use an inverted light microscope with 100X magnification to differentiate the plaque edges.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

4.11 Record results as the number of FCV plaques for each dilution of a Test Vaccine and the FCV Reference Control.

4.12 Calculate the FCV titers of the Test Vaccine and the FCV Reference Control from the dilutions with plaque counts containing 10-100 plaques. The titer may be expressed as PFU per dose of vaccine.

Example:

During the titration of a Test FCV vaccine, the plaque count obtained at dilution 10^{-3} was 65. Calculation of the titer of this vaccine is done according to the following formula:

Test Vaccine Titer = X + p, where

X = \log_{10} of plaque count

p = \log_{10} of dilution counted

In the above example:

\log_{10} of plaque count (65) = 1.8

\log_{10} of dilution counted (10^{-3}) = 3.0

FCV Test Vaccine Titer = 1.8 + 3.0 = 4.8

Adjust the titer to the recommend Test Vaccine dose as follows:

A. divide the Test Vaccine Dose by the Inoculation Dose

Test Vaccine Dose = manufacturer's recommended vaccination dose (for this test FCV vaccine, the recommended dose is 1 ml)

Inoculation Dose = amount of diluted Test Vaccine added to each well of the Test Plate (for this test FCV vaccine, the inoculation dose is 0.2 ml)

$$\frac{\underline{\underline{1 \text{ ml dose}}}}{0.2 \text{ ml inoculum}} = 5$$

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

B. calculate \log_{10} of value in **A** and add it to the **Test Vaccine Titer** as illustrated below:

$$\text{Log of } 5 = 0.7$$

$$\text{FCV Test Vaccine Titer} = 4.8 + 0.7 = 5.5$$

Therefore, the titer of the **FCV Test Vaccine** is $10^{5.5}$ PFU₅₀/ml.

4.13 One plaque represents a single infective unit (IU), whereas 1, 50% tissue culture infective dose (TCID₅₀) is statistically equivalent to a theoretical 0.69 IU. A TCID₅₀ endpoint will be 1.44 times those expressed as PFU per unit of inoculation. Therefore, to express PFU titer as TCID₅₀ titer, multiply the PFU titer by 1.44 or add 0.16 (the log of 1.44) to the \log_{10} value of the PFU titer. From the above example, the titer of the Test Vaccine is $10^{5.66}$ TCID₅₀/ml.

5. Interpretation of the test results

5.1 Validity requirements

5.1.1 The calculated titer of the FCV Reference Control must fall within plus or minus 2 standard deviations (± 2 SD) of its mean titer, as established from a minimum of 10 previously determined titers.

5.1.2 The uninoculated cell control cannot exhibit any plaques, cytopathic effects (CPE), or cloudy media that would indicate any contamination.

5.1.3 For endpoint determination, only those tenfold dilutions of the Test Vaccine and the FCV Reference Control inducing 10 to 100 plaques are included in the plaque counts.

5.2 If the validity requirements are not met, then the assay is considered a **NO TEST** and can be retested without prejudice.

5.3 If the validity requirements are met and the titer of the Test Vaccine is greater than or equal to the titer contained in the Animal and Plant Health Inspection Service (APHIS) filed Outline of Production for the product under test, the Test Vaccine is considered **SATISFACTORY**.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

5.4 In a valid test, if the titer of the Test Vaccine is less than the titer contained in the APHIS filed Outline of Production for the product under test, the Test Vaccine may be retested in accordance with the Code of Federal Regulations, Title 9, Part 113.8.b.

6. Report of test results

Report results as PFU or TCID₅₀ per dose of Test Vaccine.

7. References

7.1 Code of Federal Regulations, Title 9, Part 113.314, U.S. Government Printing Office, Washington, DC, 2004.

7.2 Cottral, GE. *Manual of standardized methods for veterinary microbiology*. Comstock Publishing Associates. Ithaca, NY, 1978, pg. 731.

7.3 Davis, Dulbecco, Eisen, Ginsberg, ed. *Microbiology including Immunology and Molecular Genetics*, 3rd ed. Harper and Row, Hagerstown, 1980, pg. 880.

8. Summary of revisions

This document was revised to clarify the practices currently in use at the Center for Veterinary Biologics and to provide additional detail. While no significant changes were made that impact the outcome of the test, the following changes were made to the document:

- **1.2** "Key Words" has been deleted.
- **2.2.3.2** The amount of sodium bicarbonate (NaHCO₂) has been changed from 2.2g to 1.1g.
- **2.2.5.8** 50 µg/ml gentamicin sulfate has been changed to 100 µg/ml and the use of penicillin and streptomycin has been deleted.
- **4.6** Ninety-six hours has been changed to 4 days ± 1 day.
- **4.12** The determination of virus titer for each Test Vaccine has been rewritten to correct errors in calculations using the example given.

Supplemental Assay Method for the Titration of Feline Calicivirus in Cell Culture

- **5.1.3** The endpoint determination of plaques has been clarified.
- The refrigeration temperatures have been changed from $4^{\circ} \pm 2^{\circ}\text{C}$ to $2^{\circ}\text{--}7^{\circ}\text{C}$. This reflects the parameters established and monitored by the Rees system.
- "Test Serial" has been changed to "Test Vaccine" throughout the document.
- "Reference Reagent Sheet" has been changed to "Reagent Data Sheet" throughout the document.
- The footnotes have been deleted with any pertinent references now noted next to the individual items.